

BARYONS IN A CHIRAL CONSTITUENT QUARK MODEL

L. Ya. Glozman

Institute for Theoretical Physics, University of Graz, Austria

Abstract

In the low-energy regime light and strange baryons should be considered as systems of constituent quarks with confining interaction and a chiral interaction that is mediated by Goldstone bosons as well as by vector and scalar mesons. The flavor-spin structure and sign of the short-range part of the spin-spin force reduces the $SU(6)_{FS}$ symmetry down to $SU(3)_F \times SU(2)_S$, induces hyperfine splittings and provides correct ordering of the lowest states with positive and negative parity. There is a cancellation of the tensor force from pseudoscalar- and vector-exchanges in baryons. The spin-orbit interactions from ρ -like and ω -like exchanges also cancel each other in baryons while they produce a big spin-orbit force in NN system. A unified description of light and strange baryon spectra calculated in a semirelativistic framework is presented. It is demonstrated that the same short-range part of spin-spin interaction between the constituent quarks induces a strong short-range repulsion in NN system when the latter is treated as $6Q$ system. Thus one can achieve a simultaneous understanding of a baryon structure and baryon-baryon interaction in the low-energy regime.

1 Introduction

Our aim in physics is not only to calculate some observable and get a correct number but mainly to understand a physical picture responsible for the given phenomenon. It very often happens that a theory formulated in terms of fundamental degrees of freedom cannot answer such a question since it becomes overcomplicated at the related scale. Thus a main task in this case is to select those degrees of freedom which are indeed essential. For instance, the fundamental degrees of freedom in crystals are ions in the lattice, electrons and the electromagnetic field. Nevertheless, in order to understand electric conductivity, heat capacity, etc. we instead work with "heavy electrons" with dynamical mass, phonons and their interaction. In this case a complicated electromagnetic interaction of the electrons with the ions in the lattice is "hidden" in the dynamical mass of the electron and the interactions among ions in the lattice are eventually responsible for the collective excitations of the lattice - phonons, which are Goldstone bosons of the spontaneously broken translational invariance in the lattice of ions. As a result, the theory becomes rather simple - only the electron and phonon degrees of freedom and their interactions are essential for all the properties of crystals mentioned above.

Quite a similar situation takes place in QCD. One hopes that sooner or later one can solve the full nonquenched QCD on the lattice and get the correct nucleon and pion mass in terms of underlying degrees of freedom: current quarks and gluon fields. However, QCD at the scale of 1 GeV becomes too complicated, and hence it is rather difficult to

say in this case what kind of physics, inherent in QCD, is relevant to the nucleon mass and its low-energy properties. In this lecture I will try to answer this question. I will show that it is the spontaneous breaking of chiral symmetry which is the most important QCD phenomenon in this case, and that beyond the scale of spontaneous breaking of chiral symmetry light and strange baryons can be viewed as systems of three constituent quarks which interact by the exchange of Goldstone bosons (pseudoscalar mesons), vector and scalar mesons (which could be considered as a representation of a correlated Goldstone boson exchange) and are subject to confinement.

2 Spontaneous Breaking of Chiral Symmetry and its Implications

At low temperatures and densities the $SU(3)_L \times SU(3)_R$ chiral symmetry of QCD Lagrangian is spontaneously broken down to $SU(3)_V$ by the QCD vacuum (in the large N_c limit it would be $U(3)_L \times U(3)_R \rightarrow U(3)_V$). A direct evidence for the spontaneously broken chiral symmetry is a nonzero value of the quark condensates for the light flavors $\langle |\bar{q}q| \rangle \approx -(240 - 250 \text{ MeV})^3$, which represent the order parameter. That this is indeed so, we know from three sources: current algebra, QCD sum rules, and lattice gauge calculations. There are two important generic consequences of the spontaneous breaking of chiral symmetry (SBCS). The first one is an appearance of the octet of pseudoscalar mesons of low mass, π, K, η , which represent the associated approximate Goldstone bosons (in the large N_c limit the flavor singlet state η' should be added). The second one is that valence (practically massless) quarks acquire a dynamical mass, which have been called historically constituent quarks. Indeed, the nonzero value of the quark condensate itself implies at the formal level that there should be rather big dynamical mass, which could be in general a moment-dependent quantity. Thus the constituent quarks should be considered as quasiparticles whose dynamical mass comes from the nonperturbative gluon and quark-antiquark dressing. The flavor-octet axial current conservation in the chiral limit tells that the constituent quarks and Goldstone bosons should be coupled with the strength $g = g_A M / f_\pi$ [1], which is a quark analog of the famous Goldberger-Treiman relation. We cannot say at the moment for sure what is the microscopical mechanism for SBCS in QCD. Any sufficiently strong scalar interaction between quarks will induce the SBCS (e.g. the instanton - induced interaction contains the scalar part, or it can be generated by monopole condensation, etc.).

All these general aspects of SBCS are well illustrated by the Nambu and Jona-Lasinio model [2], where the constituent mass is generated by the scalar part of some nonperturbative local gluonic interaction between current quarks while its pseudoscalar part gives rise to a relativistic deeply-bound pseudoscalar $Q\bar{Q}$ systems as Goldstone bosons.

Accordingly one arrives at the following interpretation of light and strange baryons in the low-energy regime. The highly nonperturbative gluodynamics gives rise to correlated quark-antiquark structures in the baryon sea (virtual mesons). At the same time the current valence quarks get dressed by the quark condensates and by the meson loops. The strongly-correlated quark-antiquark pairs in the pseudoscalar channel manifest themselves by virtual pseudoscalar mesons, while the weakly-correlated pairs in other channels - by

vector, etc mesons. When one integrates over the meson fields in the baryon wave function one arrives at the simple QQQ Fock component with confined constituent quarks and with residual interaction between them mediated by the corresponding meson fields [3].

The complimentary description of the vector-meson fields as well as of the scalar ones as arising from the correlated Goldstone bosons is also possible, which does not contradict, however, to their interpretation as weakly bound $Q\bar{Q}$ systems.

3 The Goldstone Boson Exchange Interaction

The coupling of the constituent quarks and the pseudoscalar Goldstone bosons will (in the $SU(3)_F$ symmetric approximation) have the form $g/(2m)\bar{\psi}\gamma_\mu\gamma_5\vec{\lambda}^F\cdot\psi\partial^\mu\vec{\phi}$ within the nonlinear realization of chiral symmetry (it would be $ig\bar{\psi}\gamma_5\vec{\lambda}^F\cdot\vec{\phi}\psi$ within the linear σ -model chiral symmetry representation). A coupling of this form, in a nonrelativistic reduction for the constituent quark spinors, will – to lowest order – give rise the $\sim\vec{\sigma}\cdot\vec{q}\vec{\lambda}^F$ structure of the meson-quark vertex, where $\vec{q}$ is meson momentum. Thus, the structure of the potential between quarks "i" and "j" in momentum representation is

$$V(\vec{q}) \sim \vec{\sigma}_i \cdot \vec{q} \sigma_j \cdot \vec{q} \vec{\lambda}_i^F \cdot \vec{\lambda}_j^F D(q^2) F^2(q^2), \quad (1)$$

where $D(q^2)$ is dressed Green function for chiral field which includes both nonlinear terms of chiral Lagrangian and fermion loops, $F(q^2)$ is meson-quark formfactor which takes into account the internal structure of quasiparticles. At big distances ($\vec{q} \rightarrow 0$), one has $D(q^2) \rightarrow D_0(\vec{q}^2) = -(\vec{q}^2 + \mu^2)^{-1} \neq \infty$ and $F(q^2) \rightarrow 1$. It then follows from (1) that $V(\vec{q} = 0) = 0$, which means that the volume integral of the Goldstone boson exchange (GBE) interaction should vanish,

$$\int d\vec{r} V(\vec{r}) = 0. \quad (2)$$

This sum rule is not valid in the chiral limit, however, where $\mu = 0$ and, hence, $D_0(\vec{q} = 0) = \infty$.

The sum rule (2) is trivial for the tensor component of the pseudoscalar - exchange interaction since the tensor force component automatically vanishes on averaging over the directions of $\vec{r}$. But for the spin-spin component the sum rule (2) indicates that there must be a strong short-range term. Indeed, at big interquark separations the spin-spin component of the pseudoscalar-exchange interaction is $V(r) \sim \frac{e^{-\mu r}}{r}$, it then follows from the sum rule above that at short interquark separations the spin-spin interaction should be opposite in sign as compared to the Yukawa tail and very strong:

$$H_\chi \sim - \sum_{i < j} V(\vec{r}_{ij}) \vec{\lambda}_i^F \cdot \vec{\lambda}_j^F \vec{\sigma}_i \cdot \vec{\sigma}_j. \quad (3)$$

It is this short-range part of the Goldstone boson exchange (GBE) interaction between the constituent quarks that is of crucial importance for baryons: it has a sign appropriate to reproduce the level splittings and dominates over the Yukawa tail towards short

distances. In a oversimplified consideration with a free Klein-Gordon Green function instead of the dressed one in (1) and with $F(q^2) = 1$, one obtains the following spin-spin component of $Q - Q$ interaction:

$$V(r) = \frac{g^2}{4\pi} \frac{1}{3} \frac{1}{4m_i m_j} \vec{\sigma}_i \cdot \vec{\sigma}_j \vec{\lambda}_i^F \cdot \vec{\lambda}_j^F \left\{ \mu^2 \frac{e^{-\mu r}}{r} - 4\pi \delta(\vec{r}) \right\}. \quad (4)$$

In the chiral limit only the negative short-range part of the GBE interaction survives.

4 The Vector-Meson-Exchange Interaction

As a representation of a multiple correlated Goldstone-boson-exchange one can include the exchange by a scalar flavor-singlet " σ "-meson as well as by vector mesons.

The " σ "-meson exchange does not play a principal role in baryons since it does not contain the spin-spin and tensor-force component. Its attractive central potential as well as a weak spin-orbit force can be effectively included into confining interaction in baryons. However, these forces are of significant importance in NN system at medium range.

The coupling of constituent quarks with the octet vector meson field $\vec{v}^\mu$ in the $SU(3)_F$ - symmetric approximation is

$$L^v = -g^v \bar{\psi} \gamma_\mu \vec{\lambda}^F \cdot \psi \vec{v}^\mu + \frac{g^t}{2m} \bar{\psi} \sigma_{\mu\nu} \vec{\lambda}^F \cdot \psi \partial^\nu \vec{v}^\mu, \quad (5)$$

where g^v and g^t are the vector- and tensor coupling constants. A coupling of this form - to lowest order - will give rise the spin-spin, tensor, spin-orbit, and central interactions between the constituent quarks.

It is very instructive to compare signs of the spin-spin and tensor components of pseudoscalar- and vector-exchange interactions. The spin-spin and tensor components of the pseudoscalar-exchange interaction arise from the $\vec{\sigma} \cdot \vec{\nabla}$ structure of the pseudoscalar-meson – constituent quark vertex as:

$$(\vec{\sigma}_i \cdot \vec{\nabla})(\vec{\sigma}_j \cdot \vec{\nabla}) = \frac{1}{3}(\vec{\sigma}_i \cdot \vec{\sigma}_j) \nabla^2 + \frac{1}{3} \left[3(\vec{\sigma}_i \cdot \vec{\nabla})(\vec{\sigma}_j \cdot \vec{\nabla}) - (\vec{\sigma}_i \cdot \vec{\sigma}_j) \nabla^2 \right]. \quad (6)$$

The coupling of the vector-meson to constituent quark gives $\vec{\sigma} \times \vec{\nabla}$ structure of the vertex. Therefore the spin-spin and tensor components of the vector-meson exchange interaction arise as

$$(\vec{\sigma}_i \times \vec{\nabla})(\vec{\sigma}_j \times \vec{\nabla}) = \frac{2}{3}(\vec{\sigma}_i \cdot \vec{\sigma}_j) \nabla^2 - \frac{1}{3} \left[3(\vec{\sigma}_i \cdot \vec{\nabla})(\vec{\sigma}_j \cdot \vec{\nabla}) - (\vec{\sigma}_i \cdot \vec{\sigma}_j) \nabla^2 \right]. \quad (7)$$

Comparing (6) with (7) one observes that the spin-spin component of the vector-exchange has the same sign as the spin-spin component of the pseudoscalar-exchange interaction and is "two times stronger" while their tensor components have opposite signs.

The $\vec{\sigma} \times \vec{\nabla}$ structure of the vector-meson – constituent quark vertex also suggests that the spin-spin component of the vector-meson-exchange interaction should satisfy the same sum rule (2) as the pseudoscalar-exchange interaction. Then similar to pseudoscalar-exchange interaction there should be a short-range term in the vector-meson exchange

interaction of the form (3). Again, if one neglects the spatial structure of the meson-quark vertex and uses a free Green function for the vector-meson field, this short range term is described by a δ - function piece similar to equation (4).

Summarizing, both vector- and pseudoscalar-exchange interactions produce the short-range spin-flavor force (3) while their tensor forces largely cancel each other in baryons. This observation is crucial for the present model.

The spin-orbit component associated with the vector-meson exchange interaction is big and empirically very important in NN system. So the question is why this spin-orbit force is big in NN system and becomes inessential in baryons, where the spin-orbit splittings are generally very small (see e.g. at $N(1535) - N(1520)$, $N(1650) - N(1700) - N(1675)$, ... LS multiplets). The reason for this remarkable phenomenon is an explicit flavor dependence of the vector-meson-exchange LS force.

Consider for simplicity the $SU(2)_T$ case, i.e. an exchange by ρ - and ω -mesons between U and D quarks. The ρ -meson is isovector and the ρ -meson exchange potential contains the factor $\vec{\tau}_i \cdot \vec{\tau}_j$. The ω -meson is isoscalar and the ω -exchange interaction does not contain the isospin-dependent factor. In 3P_J NN partial wave the isospin of the two-nucleon system is $T = 1$ and the isospin matrix element for the ρ -meson exchange potential is $\langle T = 1 | \vec{\tau}_i \cdot \vec{\tau}_j | T = 1 \rangle = 1$. Thus both ω - and ρ -meson exchange spin-orbit forces contribute with the same sign in NN system. Numerically the contribution from ω -exchange is 2.5-3 times bigger than from the ρ -meson exchange spin-orbit force.

Now let us consider 3P_J state of two light quarks in baryon. In the present case the isospin of the quark pair is $T = 0$, due to the presence of the color part of wave function. Then for the ρ -meson exchange one obtains $\langle T = 0 | \vec{\tau}_i \cdot \vec{\tau}_j | T = 0 \rangle = -3$. Thus the ρ -meson exchange spin-orbit force obtains opposite sign and becomes strongly enhanced in baryons. As a result one observes a strong cancellation of the ρ - and ω -meson exchange spin-orbit forces in baryons and the net weak spin-orbit interaction does not induce appreciable splittings in baryons.

Numerical details about both the tensor- and spin-orbit force cancellation in baryons can be found in ref. [4].

5 The Flavor-Spin Hyperfine Interaction and the Structure of the Baryon Spectrum

Summarizing previous sections one concludes that the pseudoscalar- and vector-meson exchange interactions produce strong flavor-spin interaction (3) at short range while the net tensor and spin-orbit forces are rather weak. That the net spin-orbit and tensor interactions between constituent quarks *in baryons* should be weak also follows from the typically small splittings in LS-multiplets, which are of the order 10-30 MeV. These small splittings should be compared with the hyperfine splittings produced by spin-spin force, which are of the order of $\Delta - N$ splitting. Thus, indeed, in baryons it is the spin-spin interaction (3) between constituent quarks is of crucial importance.

Consider first, for the purposes of illustration, a schematic model which neglects the radial dependence of the potential function $V(r)$ in (3), and assume a harmonic confine-

ment among quarks as well as $m_u = m_d = m_s$. In this model

$$H_\chi = - \sum_{i < j} C_\chi \vec{\lambda}_i^F \cdot \vec{\lambda}_j^F \vec{\sigma}_i \cdot \vec{\sigma}_j. \quad (8)$$

Note, that contrary to the color-magnetic interaction from perturbative one-gluon exchange, the chiral interaction is explicitly flavor-dependent. It is this circumstance which allows to solve the long-standing problem of ordering of the lowest positive-negative parity states.

If the only interaction between the quarks were the flavor- and spin-independent harmonic confining interaction, the baryon spectrum would be organized in multiplets of the symmetry group $SU(6)_{\text{FS}} \times U(6)_{\text{conf}}$. In this case the baryon masses would be determined solely by the orbital structure, and the spectrum would be organized in an *alternative sequence of positive and negative parity states*, i.e. in this case the spectrum would be: ground state of positive parity ($N = 0$ shell, N is the number of harmonic oscillator excitations in a 3-quark state), first excited band of negative parity ($N = 1$), second excited band of positive parity ($N = 2$), etc.

The Hamiltonian (8), within a first order perturbation theory, reduces the $SU(6)_{\text{FS}} \times U(6)_{\text{conf}}$ symmetry down to $SU(3)_F \times SU(2)_S \times U(6)_{\text{conf}}$, which automatically implies a splitting between the octet and decuplet baryons (e.g. the Δ resonance becomes heavier than nucleon).

Let us now see how the pure confinement spectrum above becomes modified when the GBE Hamiltonian (8) is switched on. For the octet states N , Λ , Σ , Ξ ($N = 0$ shell) as well as for their first radial excitations of positive parity $N(1440)$, $\Lambda(1600)$, $\Sigma(1660)$, $\Xi(?)$ ($N = 2$ shell) the expectation value of the Hamiltonian (8) is $-14C_\chi$. For the decuplet states Δ , $\Sigma(1385)$, $\Xi(1530)$, Ω ($N = 0$ shell) the corresponding matrix element is $-4C_\chi$. In the negative parity excitations ($N = 1$ shell) in the N , Λ and Σ spectra ($N(1535)$ - $N(1520)$, $\Lambda(1670)$ - $\Lambda(1690)$ and $\Sigma(1750)$ - $\Sigma(?)$) the contribution of the interaction (8) is $-2C_\chi$. The first negative parity excitation in the Λ spectrum ($N = 1$ shell) $\Lambda(1405)$ - $\Lambda(1520)$ is flavor singlet and, in this case, the corresponding matrix element is $-8C_\chi$. The latter state is unique and is absent in other spectra due to its flavor-singlet nature.

These matrix elements alone suffice to prove that the ordering of the lowest positive and negative parity states in the baryon spectrum will be correctly predicted by the chiral boson exchange interaction (8). The constant C_χ may be determined from the N - Δ splitting to be 29.3 MeV. The oscillator parameter $\hbar\omega$, which characterizes the effective confining interaction, may be determined as one half of the mass differences between the first excited $\frac{1}{2}^+$ states and the ground states of the baryons, which have the same flavor-spin, flavor and spin symmetries (e.g. $N(1440)$ - N , $\Lambda(1600)$ - Λ , $\Sigma(1660)$ - Σ), to be $\hbar\omega \simeq 250$ MeV. Thus the two free parameters of this simple model are fixed and we can make now predictions.

In the N , Λ and Σ sectors the mass difference between the lowest excited $\frac{1}{2}^+$ states ($N(1440)$, $\Lambda(1600)$, and $\Sigma(1660)$) and $\frac{1}{2}^- - \frac{3}{2}^-$ negative parity pairs ($N(1535)$ - $N(1520)$, $\Lambda(1670)$ - $\Lambda(1690)$, and $\Sigma(1750)$ - $\Sigma(?)$), respectively) will then be

$$N, \Lambda, \Sigma : \quad m(\frac{1}{2}^+) - m(\frac{1}{2}^- - \frac{3}{2}^-) = 250 \text{ MeV} - C_\chi(14 - 2) = -102 \text{ MeV}, \quad (9)$$

whereas for the lowest states in the Λ system ($\Lambda(1600)$, $\Lambda(1405)$ - $\Lambda(1520)$) it should be

$$\Lambda : \quad m(\frac{1}{2}^+) - m(\frac{1}{2}^- - \frac{3}{2}^-) = 250 \text{ MeV} - C_\chi(14 - 8) = 74 \text{ MeV}. \quad (10)$$

This simple example shows how the chiral interaction provides different ordering of the lowest positive and negative parity excited states in the spectra of the nucleon and the Λ -hyperon. This is a direct consequence of the symmetry properties of the boson-exchange interaction [3]. Namely, completely symmetric FS state in the $N(1440)$, $\Lambda(1600)$ and $\Sigma(1660)$ positive parity resonances from the $N = 2$ band feels a much stronger attractive interaction than the mixed symmetry state in the $N(1535)$ - $N(1520)$, $\Lambda(1670)$ - $\Lambda(1690)$ and $\Sigma(1750)$ - $\Sigma(?)$ resonances of negative parity ($N = 1$ shell). Consequently the masses of the positive parity states $N(1440)$, $\Lambda(1600)$ and $\Sigma(1660)$ are shifted down relative to the other ones, which explains the reversal of the otherwise expected "normal ordering". The situation is different for $\Lambda(1405)$ - $\Lambda(1520)$ and $\Lambda(1600)$, as the flavor state of $\Lambda(1405)$ - $\Lambda(1520)$ is totally antisymmetric. Because of this the $\Lambda(1405)$ - $\Lambda(1520)$ gains an attractive energy, which is comparable to that of the $\Lambda(1600)$, and thus the ordering suggested by the confining oscillator interaction is maintained.

Note that the problem of the relative position of positive-negative parity states cannot be solved with other types of hyperfine interactions between constituent quarks (the colour-magnetic and instanton-induced ones).

6 Semirelativistic Chiral Constituent Quark Model

In the semirelativistic chiral constituent quark model [5, 4] the dynamical part of the Hamiltonian consists of linear pairwise confining interaction with the string tension fixed to the known value 1 GeV/fm from Regge slopes (which also follows from the heavy quarkonium spectroscopy and lattice calculations), and the chiral interaction, mediated by pseudoscalar, scalar and vector-meson exchanges. Both the flavor-octet and flavor-singlet pseudoscalar- and vector-meson exchanges are taken into account. The coupling constants of constituent quarks with mesons are fixed from the empirically known πN , ρN , and ωN coupling constants. The "sigma-meson" - constituent quark coupling constant is taken to be equal the pion - constituent quark coupling constant, as constrained by chiral symmetry. For the constituent quark masses one takes typical values $M_{u,d} = 340$ MeV, $M_s = 500$ MeV and do not fit them. The short-range behaviour of the interaction is determined by the cut-off parameters Λ in the constituent quark - meson form-factors, which are taken in monopole form. In order to avoid a proliferation of free parameters, by assuming independent values of Λ for each meson, we adopt the linear scaling prescription $\Lambda = \Lambda_0 + \kappa\mu$, where μ is meson mass, for pseudoscalar- and vector mesons.

The kinetic-energy operator is taken in relativistic form, $H_0 = \sum_{i=1}^3 \sqrt{p_i^2 + m_i^2}$. The semirelativistic three-quark Hamiltonian was solved along the stochastic variational method [6] in momentum space. For the whole Q-Q potential the model involves a total of 4 free parameters whose numerical values are determined from the fit to all 35 confirmed low-lying states.

In fig. 1 we present the ground states as well as low-lying excited states in N , Δ , Λ ,

Σ , Ξ , and Ω spectra. From the results of fig. 1 it becomes evident that within the chiral constituent quark model a unified description of both nonstrange and strange baryon spectra is achieved in good agreement with phenomenology.

It is instructive to learn how the spin-spin interaction (3) affects the energy levels when it is switched on and its strength is gradually increased (Fig. 2). Starting out from the case with confinement only, one observes that the degeneracy of states is removed and the inversion of ordering of positive and negative parity states is achieved in the N spectrum, as well as for some states in the Λ spectrum, while the ordering of the lowest positive-negative parity states is opposite in N and Λ spectra.

It is clear that the Fock components $QQQ\pi, QQQK, \dots$ (including meson continuum) cannot be completely integrated out in favour of the meson-exchange $Q-Q$ potentials for some states above or near the corresponding meson thresholds. Such components in addition to the main one QQQ could explain e.g. an exceptionally big splitting of the flavor singlet states $\Lambda(1405) - \Lambda(1520)$, since the $\Lambda(1405)$ lies below the $\bar{K}N$ threshold and can be presented as $\bar{K}N$ bound system [7]. Note, that in the case of the present approach this old idea is completely natural and does not contradict a flavor-singlet QQQ nature of $\Lambda(1405)$, while it would be in conflict with naive constituent quark model where no room for mesons in baryons. An admixture of such components will be important in order to understand strong decays of some excited states. While technically inclusion of such components in addition to the main one QQQ in a coupled-channel approach is rather difficult task, it should be considered as one of the most important future directions.

What is an intuitive picture of the nucleon in the low-energy regime? The Goldstone bosons as well as vector meson fields couple to valence quarks. Thus the nucleon consists mostly of 3 constituent quarks which are very big objects due to their meson clouds (see fig. 3). These constituent quarks are all the time in strong overlap inside the nucleon. That is why the short-range part of chiral interaction (which is represented by the contact term in the oversimplified representation) is so crucially important inside baryons. When constituent quarks are well separated and there is a phase space for meson propagation, the long-range Yukawa tails of meson-exchange interactions become very important. It is these parts of meson exchange which produce the necessary long- and intermediate-range attraction in two-nucleon system.

7 The Baryon-Baryon Interaction in a Chiral Constituent Quark Model

If one ascribes a short-range central repulsion in the NN system to the central part of ω exchange between nucleons, then one should increase the ωN coupling constant by factor 3 as compared to its empirical value, as it is usually done in phenomenological one-boson-exchange NN potentials. Evidently the short-range repulsion in the NN system should be connected with the nucleon structure in the low-energy regime and within the quark picture should be related to fermi-nature of constituent quarks and to the specific interactions between them.

So far, all studies of the short-range NN interaction within the constituent quark model were based on the one-gluon exchange interaction between quarks. They explained




Figure 1: Energy levels of the lowest light and strange baryon states (below 1850 MeV) with total angular momentum and parity J^P . The shadowed boxes represent the experimental values with their uncertainties.




Figure 2: Level shifts of some lowest baryons as a function of the strength of the chiral interaction. Solid and dashed lines correspond to positive- and negative-parity states, respectively.




Figure 3: Nucleon as it is seen in the low-energy and low-resolution regime.

the short-range repulsion in the NN system as due to the colour-magnetic part of OGE combined with quark interchanges between 3Q clusters [8]. It has been shown, however, that there is practically no room for colour-magnetic interaction in light baryon spectroscopy and any appreciable amount of colour-magnetic interaction, in addition to chiral interaction, destroys the spectrum [9]. This conclusion is confirmed by recent lattice QCD calculations [10]. If so, the question arises which interquark interaction is responsible for the short-range NN repulsion. Below I show that the same short-range part of chiral interaction (3) which causes e.g. $N - \Delta$ splitting and produces good baryon spectra, also induces a short-range repulsion in NN system when the latter is treated as 6Q system [12].

At present one can use only a simple nonrelativistic s^3 ansatz for the nucleon wave function when one applies the quark model to NN interaction. Thus one needs first an effective nonrelativistic parametrization of chiral interaction interaction which would provide correct nucleon mass, $N - \Delta$ splitting and the nucleon stability with this ansatz. For that one can use the nonrelativistic parametrization [11] which satisfies approximately the conditions above.

In order to have a qualitative insight into the NN interaction it is convenient to use an adiabatic Born-Oppenheimer approximation for the internucleon potential:

$$V_{NN}(R) = \langle H \rangle_R - \langle H \rangle_\infty, \quad (11)$$

where R is a collective (generator) coordinate which is the separation distance between the two wells (it should not be mixed with the relative motion Jacobi coordinate), $\langle H \rangle_R$ is the lowest expectation value of the 6Q Hamiltonian at fixed R , and $\langle H \rangle_\infty$ is a mass of two well-separated nucleons ($2m_N$) calculated with the same Hamiltonian.

At the moment we are interested in what is the NN interaction at zero separation between nucleons. It has been proved by Harvey that when $R \rightarrow 0$, then in both 3S_1 and 1S_0 partial waves in the NN system only two types of orbital 6Q configurations survive [13]: $|s^6[6]_O\rangle$ and $|s^4p^2[42]_O\rangle$, where $[f]_O$ is Young diagram, describing spatial permutational symmetry in 6Q system. There are a few different flavor-spin symmetries, compatible with the spatial symmetries above: $[6]_O[33]_{FS}$, $[42]_O[33]_{FS}$, $[42]_O[51]_{FS}$, $[42]_O[411]_{FS}$, $[42]_O[321]_{FS}$, and $[42]_O[2211]_{FS}$. Thus, in order to evaluate the NN interaction at zero separation between nucleons it is necessary to diagonalize a 6Q Hamiltonian in the basis above and use the procedure (11).

From the adiabatic Born-Oppenheimer approximation (11) we find that $V_{NN}(R=0)$ is highly repulsive in both 3S_1 and 1S_0 partial waves, with the core being of order 1 GeV. This repulsion implies a strong suppression of the NN wave function in the nucleon overlap region.

Due to the specific flavor-spin symmetry of chiral interaction between quarks the configuration $s^4p^2[42]_O[51]_{FS}$ becomes highly dominant among other possible 6Q configurations at zero separation between nucleons (however, the "energy" of this configuration is much higher than the energy of two well-separated nucleons, that is why there is a strong short-range repulsion in NN system). The symmetry structure of this dominant configuration induces "an additional" effective repulsion, related to the "Pauli forbidden state" in this case, and the s-wave NN relative motion wave function has a node at short range [14]. The existence of a strong repulsion, related to the energy ballance, discussed

above, suggests, however, that the amplitude of the oscillating NN wave function at short range will be strongly suppressed.

Thus, within the chiral constituent quark model one has all the necessary ingredients to understand microscopically the NN interaction. There appears strong short-range repulsion from the same short-range part of chiral interaction which also produces hyperfine splittings in baryon spectroscopy. The long- and intermediate-range attraction in the NN system is automatically implied by the Yukawa part of pion-exchange and correlated two-pion exchanges (σ -exchange) between quarks belonging to different nucleons. The necessary tensor and spin-orbit force comes from the pseudoscalar- and vector-exchange Yukawa tails.

What will be a short-range interaction in other YN and YY systems? In the chiral limit there is no difference between all octet baryons: N, Λ, Σ, Ξ . Thus if one explains a strong short-range repulsion in NN system as related mostly to the spontaneous breaking of chiral symmetry, as above, then the same short-range repulsion should persist in other YN and YY systems [15]. Of course, due to the explicit chiral symmetry breaking the strength of this repulsion should be essentially different as compared to that one in NN system. One can naively expect that it will be weaker.

Acknowledgement

It is my pleasure to thank D.O.Riska, Z.Papp, W.Plessas, K.Varga, R. Wagenbrunn, Fl. Stancu, and S. Pepin, in collaboration with whom different results discussed in this talk have been obtained. I also thank organizers of the school for the invitation to give this lecture.

References

- [1] A. Manohar and H. Georgi, Nucl. Phys. **B234** (1984) 189; D. I. Diakonov and V. Yu. Petrov, Nucl. Phys. **B272** (1986) 457.
- [2] Y. Nambu and G. Jona-Lasinio, Phys. Rev. **122** (1961) 345; U. Vogl and W. Weise, Progr. Part. Nucl. Phys. **27** (1991) 195; S. P. Klevansky, Rev. Mod. Phys. **64** (1992) 649; T. Hatsuda and T. Kunihiro, Phys. Rep. **247** (1994) 221.
- [3] L. Ya. Glozman and D. O. Riska, Physics Reports **268** (1996) 263.
- [4] L. Ya. Glozman, W. Plessas, K. Varga, R. Wagenbrunn, in preparation.
- [5] L. Ya. Glozman, W. Plessas, K. Varga, R. Wagenbrunn, hep-ph/9706507.
- [6] K. Varga and Y. Suzuki, Phys. Rev. **C52** (1995) 2885; K. Varga, Y. Ohbayasi, and Y. Suzuki, Phys. Lett. **B396** (1997) 1.
- [7] R. H. Dalitz and A. Deloff, J. Phys. **G 17** (1991) 289 (and references therein).
- [8] M. Oka and K. Yazaki, Int. Rev. Nuc. Phys., vol. 1 (Quarks and Nuclei, ed. W. Weise), World Scientific, Singapore, p. 490 (1984).

- [9] L. Ya. Glozman, Z. Papp, W. Plessas, K. Varga, and R. Wagenbrunn, Phys. Rev. **C57** (1998) [nucl-th/9705011].
- [10] K. F. Liu and S. J. Dong - paper in preparation (for a preliminary result see hep-lat/9411067).
- [11] L. Ya. Glozman, Z. Papp, and W. Plessas, Phys. Lett. **B381** (1996) 311.
- [12] Fl. Stancu, S. Pepin, and L. Ya. Glozman, Phys. Rev. **C56** (1997) 2779.
- [13] M. Harvey, Nucl. Phys. **A352** (1981) 301.
- [14] V. G. Neudatchin, Yu. F. Smirnov, and R. Tamagaki, Progr. Theor. Phys. **58** (1977) 1072.
- [15] Fl. Stancu, S. Pepin, and L. Ya. Glozman, Phys. Rev. **D57** (1998) 4393.